

## PATIENT: XXXXXXXXXXXXXXXXXX

TEST NUMBER: T-NL-XXXXX (XXXXXXXXXX)

GENDER: XYZ XX

COLLECTED: XX/XX/XXXX RECEIVED: XX/XX/XXXX

TESTED:

XX/XX/XXXX

TEST REF: TST-NL-XXXX

PRACTITIONER XXXXXXXXXXXX

XXXXXXXXXXXXXXXXXXXX

patient's

## **TEST NAME: Urine Halides (Pre & Post Load)**

## Urine Halides; Pre & Post Loading

**lodine** μg/mg cr mg/24 hr Reference Range lodine levels include iodine and iodide 0.27 Sample 1 PRE 0.1- 0.45 μg/mg cr oxidized to iodine. Excretion percentage is calculated by dividing the mg/24hour lodine result by the lodine/lodide Sample 2 POST 25 30 0.1- 0.45 mg/24 hr dosage (in mg) recorded on the requisition form, then multiplying by 100. 60% % Excretion/24 hr

**Bromine** μg/mg cr mg/24 hr Reference Range Sample 1 PRE 3.2 7 μg/mg cr 7 mg/24 hr Sample 2 POST

**Bromine** levels represent total bromine plus bromide, as measured by ICP-MS. Bromide is antagonistic to iodide, and is abundant in commercially produced baked goods, soft drinks, pesticides, brominated chemicals and some medications.

**Fluoride** μg/mL mg/24 hr Reference Range Sample 1 PRE 0.33 1.1 μg/mL Sample 2 POST 0.29 0.78 1.3 mg/24 hr

Fluoride in urine is measured using an ion specific electrode. Fluoride is neurotoxic, compromises integrity of bone, and interferes with iodide metabolism. Primary sources of fluoride include fluoridated water, toothpaste/mouth beverages, washes, dental treatments and some medications.

Result Reference Range Creatinine 30- 225 mg/dL Sample 1 PRE 57.6 Sample 2 POST 1200 600- 2100 mg/24hr Urine Creatinine is used to account for urinary dilution effects in less than 24-hour collections and to assess the collection completeness in 24-hour collections. For estimation of glomerular filtration rate (GFR), Creatinine Clearance test is recommended.

Comments:

#1 Date Collected: 01/04/2019 #2 Date Collected:

01/05/2019 Date Received:

01/12/2019

#1 Collection Period: Random

#2 Collection Period: 24 Hr/Coll #2 Volume: 2700 ml #2 Loading Dosage: 50 MG

Date Reported: 01/18/2019 <dl: less than detection limit Method: I, Br by ICP-MS/ F by ISE Creatinine by Jaffe method

Reference ranges are representative of a healthy population under non-challenge or non-loading conditions. V04.07

Nordic Laboratories Aps

Nygade 6, 3.sal • 1164 Copenhagen K • Denmark Tlf. +45 33 75 10 00

**UK Office:** 

11 Old Factory Buildings • Stonegate • E. Sussex TN5 7DU • UK Tel: +44 (0)1580 201 687

Page 1 of 1 www.nordic-labs.com info@nordic-labs.com